

Spatial and Temporal Trends in PCB Concentrations in Fish Tissue in the Mainstem of the Delaware River




Delaware Estuary Science
and Environmental
Summit

January 26, 2015


Thomas J. Fikslin, Ph.D. and Gregory Cavallo, P.G.
Modeling, Monitoring and Assessment Branch



Presentation Themes

- ✓ Background
 - Program objectives
 - PCB TMDLs Implementation
 - Expectations
 - ✓ Sampling Design
 - ✓ Results
 - ✓ Summary
- 
- A stylized, dark teal silhouette of a mountain range is positioned in the bottom right corner of the slide, extending from the right edge towards the center.


Background

- ❑ Why monitor fish?
 - Integrator of exposure.
 - Endpoint for evaluating human health impacts from fish consumption.
 - Interstate waters.
 - Needed for Integrated Assessment of water segments by states and DRBC.
 - Funding for programs is limited.
 - Coordination w/ State partners.
- 
- A stylized, dark teal silhouette of a mountain range is positioned in the bottom right corner of the slide, partially overlapping the text area.

PCB TMDLs Implementation

- ◆ Goal is to achieve the recently adopted uniform water quality criterion of 16 picograms/Liter and eliminate the need for fish consumption advisories for Zones 2 through 6.
- ◆ For PCBs, however, contaminated sediments will continue to bleed PCBs to the water column even if all active sources of PCBs are eliminated, resulting in a significant lag time to reach the water quality criterion.

Projected, averaged Total PCB concentrations for Zones 2 & 3
Boundaries @16pg/L; current sediment condition; 99.97% all load reduction



PCB TMDLs Implementation

- ◆ TMDLs for certain hydrophobic pollutants like PCBs will require long-term strategies to achieve their goals.
- ◆ **Expectation:**

Ambient water and fish tissue concentrations will not decrease until sources of PCBs are reduced *and* sediment concentrations decline as uncontaminated sediment is deposited in the estuary.

Sampling Design

- Design considerations:
 - Locations – tidal vs. non-tidal?
 - Species – resident or migratory?
 - Analytical parameters?



Sampling Design

- ❑ Fish samples are collected from 8 sites in both the tidal (5 sites) and non-tidal (3 sites) portions of the Delaware River.
- ❑ Frequency: Yearly 2000 - 2007, 2010, 2012
- ❑ Two species of fish are collected at each site representing resident benthic and pelagic trophic levels.
 - Tidal species: white perch, channel catfish
 - Non-tidal species: smallmouth bass, white sucker
- ❑ Samples are collected by electrofishing or hook & line.


Sampling Locations 2004 to 2007, 2010, 2012

Non-Tidal Locations

Narrowsburg, NY	RM 290*
Milford, PA	RM 246
Easton, PA	RM 183
Lambertville, NJ	RM 149

Tidal Locations


Crosswicks Creek	RM 128
Tacony-Palymra Br.	RM 107
Woodbury Creek	RM 91
Raccoon Creek	RM 80
Salem River	RM 58




Analytical Methods

- ❑ Samples are composites of standard fillets, and consist of a composite of 4 to 5 fish of similar size and weight.
- ❑ Analytical Parameters & Methods:
 - Starting in 2004, all analyses were conducted by Axys Analytical LTD using Method 1668A.
 - Target analytes: all 209 PCB compounds (i.e., PCB congeners)


PCBs in Fish Tissue Delaware River Estuary 2000 to 2012




PCBs in Fish Tissue Non-Tidal Delaware River 1991 to 2012



Historical Trend in Total PCBs in Fish Tissue Tacony-Palmyra Bridge - Delaware Estuary



Historical Trend in Total PCBs in Fish Tissue Raccoon Creek - Delaware Estuary



Summary

- ◆ Since the development of the PCB TMDLs in 2003, the DRBC has analyzed fish tissue samples from 9 locations in the non-tidal and tidal portions of the Delaware River.
- ◆ Highest concentrations are observed in the urban areas of the estuary.
- ◆ Concentrations in samples collected in the non-tidal portion of the river had significantly lower concentrations of PCBs.
- ◆ PCB concentrations were higher in benthic species compared to pelagic species tested at all locations.

Summary

- ◆ As expected, tissue concentrations are not declining despite a 46% reduction in the loadings of PCBs from point sources from 2005 to 2011.
- ◆ Declines in fish tissue concentrations can be expected sooner in areas where there is less sediment contamination such as Zone 2 and Delaware Bay.
- ◆ Continued implementation of the long-term strategy developed by the co-regulators is needed to achieve the goal of eliminating the need for fish consumption advisories.

Contact Information:

thomas.fikslin@drbc.state.nj.us
(609) 477-7253

Information on the TMDLs, model development, sampling and analytical information, and other implementation requirements and resources are available on the DRBC website at:

<http://www.state.nj.us/drbc/quality/toxics/pcbs/>

